

## Wildlife Interpretations

### **General**

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

Soils are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife. It can also be used in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat.

### **Suitability Ratings**

**GOOD** - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

**FAIR** - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

**POOR** - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

### **Description of Wildlife Habitat Elements**

Grain and seed crops are seed-producing annuals used by wildlife. Examples are corn, wheat, rye, oats, barley, millet, buckwheat, and sunflowers.

Grasses and legumes are domestic perennial grasses and herbaceous legumes that are planted for wildlife food and cover. Examples are fescue, bluegrass, brome grass, timothy, orchard grass, clover, alfalfa, trefoil, reed canary grass, and crown vetch.

Wild herbaceous plants are native and naturally established herbaceous grasses, forbs, and weeds, that provide food and cover for wildlife. Examples are bluestem, indiagrass, blueberry, goldenrod, lambsquarters, dandelions, blackberry, ragweed, wheatgrass, fescue, and nightshade.

Hardwood trees and the associated woody understory provide cover for wildlife and produce nuts or other fruit, buds, catkins, twigs, bark, or foliage that wildlife eat. Examples of native plants are oak, poplar, cherry, apple, birch, beech, maple, hickory, hazelnut, black walnut, and viburnum. Examples of fruit-bearing shrubs that are

commercially available and suitable for planting on soils rated good are hawthorn, honeysuckle, sumac, silky dogwood, highbush cranberry, autumn olive, and crabapple.

Coniferous plants are cone-bearing trees, shrubs, or ground cover that furnish habitat or supply food in the form of browse, seed, or fruit-like cones. Examples are pine, spruce, hemlock, fir, yew, cedar, larch, and juniper.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites, exclusive of submerged or floating aquatics. They produce food or cover for wildlife that use wetland as habitat. Examples of wetland plants are smartweed, wild millet, rushes, sedges, reeds, wildrice, arrowhead, water plantain, pickerelweed, and cattail.

Shallow water are bodies of surface water that have an average depth of less than 5 feet and are useful as habitat for wildlife. They can be naturally wet areas, or they can be created by dams or levees, or by water-control devices in marshes or streams. Examples are muskrat marshes, waterfowl feeding areas, wildlife watering developments, beaver ponds, and other wildlife ponds.

### ***Kinds of Wildlife Habitat***

Open land habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and woodchuck.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, woodcock, thrushes, woodpeckers, owl, tree squirrels, gray fox, porcupine, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, muskrat, otter, mink, and beaver.

*This subsection includes:*

- **(a) Wildlife Management Ratings (one or two tables)**

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
1: Adler-----	Moderately limited ~flooding (moderately limited) ~wetness (slightly limited)	0.60  0.28	Moderately limited ~flooding (moderately limited) ~wetness (slightly limited)	0.60  0.28	Slightly limited ~wetness (slightly limited) ~flooding (prolonged) (slightly limited)	0.28  0.20	Slightly limited ~wetness (slightly limited) ~flooding (prolonged) (slightly limited)	0.28  0.20	Moderately limited ~wetness (moderately limited) ~flooding (prolonged) (slightly limited)	0.45  0.20
2: Amagon-----	Moderately limited ~flooding (moderately limited) ~wetness (moderately limited) ~percs slowly (moderately limited)	0.60  0.60 0.39	Moderately limited ~flooding (moderately limited) ~wetness (moderately limited) ~percs slowly (moderately limited)	0.60  0.60 0.39	Moderately limited ~wetness (moderately limited)	0.60	Moderately limited ~wetness (moderately limited)	0.60	Limited ~wetness (limited)	0.99
3B: Elk-----	Moderately limited ~flooding (moderately limited)	0.60	Moderately limited ~flooding (moderately limited)	0.60	Not limited		Not limited		Not limited	
4D: Bardley-----	Very limited ~droughty (very limited) ~high erodibility (limited) ~small stones (moderately limited)	1.00  0.80 0.46	Limited ~high erodibility (limited) ~small stones (moderately limited) ~depth to bedrock (moderately limited)	0.80  0.46 0.46	Moderately limited ~droughty (moderately limited) ~small stones (slightly limited)	0.37  0.08	Moderately limited ~depth to bedrock (moderately limited) ~droughty (moderately limited)	0.46  0.37	Moderately limited ~depth to bedrock (moderately limited) ~droughty (moderately limited)	0.46  0.37
5B: Bosket-----	Slightly limited ~droughty (slightly limited)	0.06	Not limited		Not limited		Not limited		Not limited	

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
5C: Bosket-----	Moderately limited ~moderate erodibility (moderately limited) ~droughty (slightly limited)	0.50 0.06	Moderately limited ~moderate erodibility (moderately limited)	0.50	Not limited		Not limited		Not limited	
6C2: Bosket-----	Moderately limited ~moderate erodibility (moderately limited) ~droughty (slightly limited)	0.50 0.01	Moderately limited ~moderate erodibility (moderately limited)	0.50	Not limited		Not limited		Not limited	
7: Calhoun-----	Limited ~wetness (limited) ~flooding (moderately limited) ~percs slowly (moderately limited)	0.99 0.60 0.39	Limited ~wetness (limited) ~flooding (moderately limited) ~percs slowly (moderately limited)	0.99 0.60 0.39	Limited ~wetness (limited) ~flooding (prolonged) (slightly limited)	0.99 0.20	Limited ~wetness (limited) ~flooding (prolonged) (slightly limited)	0.99 0.20	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20
8B: Captina-----	Moderately limited ~moderate erodibility (moderately limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.50 0.39 0.28	Moderately limited ~moderate erodibility (moderately limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.50 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
8B2: Captina-----	Moderately limited ~moderate erodibility (moderately limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.50 0.39 0.28	Moderately limited ~moderate erodibility (moderately limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.50 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
8C: Captina-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
8C2: Captina-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
9C: Clarksville----	Very limited ~droughty (very limited) ~small stones (very limited) ~moderate erodibility (moderately limited)	1.00 1.00 0.50	Very limited ~small stones (very limited) ~moderate erodibility (moderately limited) ~droughty (slightly limited)	1.00 0.50 0.16	Moderately limited ~small stones (moderately limited) ~droughty (slightly limited)	0.52 0.16	Moderately limited ~small stones (moderately limited) ~droughty (slightly limited)	0.47 0.16	Slightly limited ~droughty (slightly limited)	0.16

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
9D: Clarksville----	Very limited ~droughty (very limited) ~small stones (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~small stones (very limited) ~high erodibility (limited) ~droughty (slightly limited)	1.00  0.80  0.16	Moderately limited ~small stones (moderately limited) ~droughty (slightly limited)	0.52  0.16	Moderately limited ~small stones (moderately limited) ~droughty (slightly limited)	0.47  0.16	Slightly limited ~droughty (slightly limited)	0.16
9F: Clarksville----	Very limited ~droughty (very limited) ~small stones (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~small stones (very limited) ~high erodibility (limited) ~slope (moderately limited)	1.00  0.80  0.42	Moderately limited ~small stones (moderately limited) ~droughty (slightly limited)	0.52  0.16	Moderately limited ~small stones (moderately limited) ~droughty (slightly limited)	0.47  0.16	Slightly limited ~droughty (slightly limited)	0.16
10D: Clarksville----	Very limited ~droughty (very limited) ~small stones (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~small stones (very limited) ~high erodibility (limited) ~droughty (moderately limited)	1.00  0.80  0.51	Moderately limited ~droughty (moderately limited) ~small stones (slightly limited)	0.51  0.24	Moderately limited ~droughty (moderately limited) ~small stones (slightly limited)	0.51  0.01	Moderately limited ~droughty (moderately limited)	0.51
10F: Clarksville----	Very limited ~droughty (very limited) ~small stones (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~small stones (very limited) ~high erodibility (limited) ~droughty (moderately limited)	1.00  0.80  0.51	Moderately limited ~droughty (moderately limited) ~small stones (slightly limited)	0.51  0.24	Moderately limited ~droughty (moderately limited) ~small stones (slightly limited)	0.51  0.01	Moderately limited ~droughty (moderately limited)	0.51

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
11: Crowley-----	Very limited ~percs slowly (very limited) ~wetness (limited)	1.00  0.99	Very limited ~percs slowly (very limited) ~wetness (limited)	1.00  0.99	Limited ~wetness (limited)	0.99	Limited ~wetness (limited)	0.99	Very limited ~wetness (very limited)	1.00
12C: Doniphan-----	Limited ~droughty (very limited) ~small stones (moderately limited) ~moderate erodibility (moderately limited)	1.00  0.60 0.50	Moderately limited ~small stones (moderately limited) ~moderate erodibility (moderately limited) ~large stones (slightly limited)	0.60  0.50 0.09	Slightly limited ~small stones (slightly limited)	0.12	Not limited		Not limited	
12D: Doniphan-----	Limited ~droughty (very limited) ~high erodibility (limited) ~small stones (moderately limited)	1.00  0.80 0.60	Limited ~high erodibility (limited) ~small stones (moderately limited) ~large stones (slightly limited)	0.80  0.60 0.09	Slightly limited ~small stones (slightly limited)	0.12	Not limited		Not limited	
12F: Doniphan-----	Limited ~droughty (very limited) ~high erodibility (limited) ~slope (moderately limited)	1.00  0.80 0.60	Limited ~high erodibility (limited) ~slope (moderately limited) ~small stones (moderately limited)	0.80  0.60 0.60	Slightly limited ~small stones (slightly limited)	0.12	Not limited		Not limited	

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
13D: Doniphan-----	Very limited ~droughty (very limited) ~high erodibility (limited)	1.00  0.80	Limited ~high erodibility (limited) ~droughty (slightly limited)	0.80  0.14	Slightly limited ~droughty (slightly limited)	0.14	Slightly limited ~droughty (slightly limited)	0.14	Slightly limited ~droughty (slightly limited)	0.14
14B: Dubbs-----	Moderately limited ~moderate erodibility (moderately limited)	0.50	Moderately limited ~moderate erodibility (moderately limited)	0.50	Not limited		Not limited		Not limited	
15: Foley-----	Very limited ~wetness (very limited) ~percs slowly (very limited) ~moderate erodibility (moderately limited)	1.00  1.00  0.50	Very limited ~wetness (very limited) ~percs slowly (very limited) ~moderate erodibility (moderately limited)	1.00  1.00  0.50	Very limited ~wetness (very limited)	1.00	Very limited ~wetness (very limited)	1.00	Very limited ~wetness (very limited)	1.00
16D: Gasconade-----	Very limited ~droughty (very limited) ~shallow to bedrock (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~droughty (very limited) ~shallow to bedrock (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~droughty (very limited) ~too clayey (moderately limited) ~large stones (moderately limited)	1.00  0.38  0.30	Very limited ~droughty (very limited) ~shallow to bedrock (very limited) ~too clayey (moderately limited)	1.00  1.00  0.38	Very limited ~shallow to bedrock (very limited) ~droughty (very limited) ~large stones (moderately limited)	1.00  1.00  0.30
Rock outcrop---	Not rated		Not rated		Not rated		Not rated		Not rated	



## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
16F: Gasconade-----	Very limited ~droughty (very limited) ~shallow to bedrock (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~droughty (very limited) ~shallow to bedrock (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~droughty (very limited) ~too clayey (moderately limited) ~large stones (moderately limited)	1.00  0.38  0.30	Very limited ~droughty (very limited) ~shallow to bedrock (very limited) ~too clayey (moderately limited)	1.00  1.00  0.38	Very limited ~shallow to bedrock (very limited) ~droughty (very limited) ~large stones (moderately limited)	1.00  1.00  0.30
Rock outcrop---	Not rated		Not rated		Not rated		Not rated		Not rated	
17C: Gatewood-----	Limited ~droughty (very limited) ~high erodibility (limited) ~percs slowly (moderately limited)	1.00  0.80  0.39	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (moderately limited)	0.80  0.39  0.36	Moderately limited ~wetness (moderately limited)	0.36	Moderately limited ~wetness (moderately limited) ~depth to bedrock (slightly limited)	0.36  0.13	Moderately limited ~wetness (moderately limited) ~depth to bedrock (slightly limited)	0.51  0.13
18B: Hartville-----	Moderately limited ~flooding (moderately limited) ~moderate erodibility (moderately limited) ~percs slowly (moderately limited)	0.60  0.50  0.39	Moderately limited ~flooding (moderately limited) ~moderate erodibility (moderately limited) ~percs slowly (moderately limited)	0.60  0.50  0.39	Moderately limited ~wetness (moderately limited)	0.36	Moderately limited ~wetness (moderately limited)	0.36	Moderately limited ~wetness (moderately limited)	0.51

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
19: Hontas-----	Moderately limited ~flooding (moderately limited) ~wetness (moderately limited)	0.60  0.36	Moderately limited ~flooding (moderately limited) ~wetness (moderately limited)	0.60  0.36	Moderately limited ~wetness (moderately limited)	0.36	Moderately limited ~wetness (moderately limited)	0.36	Moderately limited ~wetness (moderately limited)	0.51
20: Houlka-----	Very limited ~percs slowly (very limited) ~flooding (moderately limited) ~wetness (moderately limited)	1.00  0.60  0.60	Very limited ~percs slowly (very limited) ~flooding (moderately limited) ~wetness (moderately limited)	1.00  0.60  0.60	Moderately limited ~wetness (moderately limited) ~flooding (prolonged) (slightly limited) ~too clayey (slightly limited)	0.60  0.20  0.16	Moderately limited ~wetness (moderately limited) ~flooding (prolonged) (slightly limited) ~too clayey (slightly limited)	0.60  0.20  0.16	Limited ~wetness (limited) ~flooding (prolonged) (slightly limited)	0.99  0.20
21: Kobel-----	Very limited ~wetness (very limited) ~percs slowly (very limited) ~droughty (limited)	1.00  1.00  0.63	Very limited ~wetness (very limited) ~percs slowly (very limited) ~too clayey (limited)	1.00  1.00  0.61	Very limited ~wetness (very limited) ~flooding (prolonged) (limited) ~too clayey (limited)	1.00  0.90  0.61	Very limited ~wetness (very limited) ~flooding (prolonged) (limited) ~too clayey (limited)	1.00  0.90  0.61	Very limited ~wetness (very limited) ~flooding (prolonged) (limited)	1.00  0.90
22: Lafe-----	Very limited ~wetness (very limited) ~percs slowly (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~wetness (very limited) ~percs slowly (very limited) ~high erodibility (limited)	1.00  1.00  0.80	Very limited ~wetness (very limited)	1.00	Very limited ~wetness (very limited)	1.00	Very limited ~wetness (very limited)	1.00

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
23B: Loring-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
23B2: Loring-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
23C: Loring-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
23C2: Loring-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
23D: Loring-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
23D2: Loring-----	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Limited ~high erodibility (limited) ~percs slowly (moderately limited) ~wetness (slightly limited)	0.80 0.39 0.28	Slightly limited ~wetness (slightly limited)	0.28	Slightly limited ~wetness (slightly limited)	0.28	Moderately limited ~wetness (moderately limited)	0.45
24A: Midco-----	Very limited ~droughty (very limited) ~flooding (moderately limited) ~small stones (moderately limited)	1.00 0.60 0.30	Moderately limited ~flooding (moderately limited) ~small stones (moderately limited) ~droughty (slightly limited)	0.60 0.30 0.18	Slightly limited ~droughty (slightly limited) ~small stones (slightly limited)	0.18 0.03	Slightly limited ~droughty (slightly limited)	0.18	Slightly limited ~droughty (slightly limited)	0.18
25: Nolin-----	Moderately limited ~flooding (moderately limited)	0.60	Moderately limited ~flooding (moderately limited)	0.60	Slightly limited ~flooding (prolonged) (slightly limited)	0.20	Slightly limited ~flooding (prolonged) (slightly limited)	0.20	Slightly limited ~flooding (prolonged) (slightly limited)	0.20
26B: Peridge-----	Moderately limited ~moderate erodibility (moderately limited)	0.50	Moderately limited ~moderate erodibility (moderately limited)	0.50	Not limited		Not limited		Not limited	

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
26C: Peridge-----	Limited ~high erodibility (limited)	0.80	Limited ~high erodibility (limited)	0.80	Not limited		Not limited		Not limited	
26C2: Peridge-----	Limited ~high erodibility (limited)	0.80	Limited ~high erodibility (limited)	0.80	Not limited		Not limited		Not limited	
27: Secesh-----	Limited ~droughty (limited) ~flooding (moderately limited)	0.69 0.60	Moderately limited ~flooding (moderately limited)	0.60	Not limited		Not limited		Not limited	
28: Tuckerman-----	Very limited ~wetness (very limited) ~flooding (moderately limited) ~percs slowly (slightly limited)	1.00 0.60 0.13	Very limited ~wetness (very limited) ~flooding (moderately limited) ~percs slowly (slightly limited)	1.00 0.60 0.13	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20
29B: Tuckerman-----	Very limited ~wetness (very limited) ~flooding (moderately limited) ~moderate erodibility (moderately limited)	1.00 0.60 0.50	Very limited ~wetness (very limited) ~flooding (moderately limited) ~moderate erodibility (moderately limited)	1.00 0.60 0.50	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20	Very limited ~wetness (very limited) ~flooding (prolonged) (slightly limited)	1.00 0.20

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
29B: (cont)										
Bosket-----	Moderately limited ~moderate erodibility (moderately limited)	0.50	Moderately limited ~moderate erodibility (moderately limited)	0.50	Not limited		Not limited		Not limited	
	~droughty (slightly limited)	0.06								
30:										
Wideman-----	Limited ~droughty (very limited)	1.00	Moderately limited ~flooding (moderately limited)	0.60	Moderately limited ~too sandy (moderately limited)	0.50	Moderately limited ~too sandy (moderately limited)	0.50	Not limited	
	~flooding (moderately limited)	0.60	~too sandy (moderately limited)	0.50						
	~too sandy (moderately limited)	0.50								
31B:										
Wilderness-----	Very limited ~droughty (very limited)	1.00	Very limited ~droughty (very limited)	1.00	Very limited ~droughty (very limited)	1.00	Very limited ~droughty (very limited)	1.00	Very limited ~droughty (very limited)	1.00
	~small stones (limited)	0.65	~small stones (limited)	0.65	~wetness (moderately limited)	0.60	~wetness (moderately limited)	0.60	~wetness (limited)	0.99
	~wetness (moderately limited)	0.60	~wetness (moderately limited)	0.60	~small stones (slightly limited)	0.13				
32:										
Pits-----	Not rated		Not rated		Not rated		Not rated		Not rated	

## Butler County and Part of Ripley County, Missouri: Detailed Soil Map Legend

## Wildlife Habitat

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

Map symbol and soil name	Grain and seed crops (for use as food and cover)		Domestic grasses and legumes (for use as food and cover)		Upland wild herbaceous plants		Upland shrubs and vines		Upland deciduous trees	
	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value	Rating class and limiting features	Value
33: Calhoun-----	Limited ~wetness (limited) ~high erodibility (limited) ~percs slowly (moderately limited)	0.99  0.80  0.39	Limited ~wetness (limited) ~high erodibility (limited) ~percs slowly (moderately limited)	0.99  0.80  0.39	Limited ~wetness (limited)	0.99	Limited ~wetness (limited)	0.99	Very limited ~wetness (very limited)	1.00
W: Water-----	Not rated		Not rated		Not rated		Not rated		Not rated	